


(NASA-CR-197882) VOYAGER TO SATURN, VOLUME 4 (JPL) 87 P N95-70915


Unclas

29/91 0043479






Assembled in this book are pictures of Saturn, its satellites, and its rings as taken by the Voyager spacecraft. Never before in the history of the space age has a single mission produced such an abundance of knowledge about the outer planets of our solar system. If you do not believe the old adage that "a picture's worth a thousand words," look within to discover the treasures unlocked by Voyager's encounter with Saturn.




### TABLE OF CONTENTS

HQ#	JPL#	Description
80-HC-90 80-H-117		Voyager 1 194,400,000 miles from Saturn
80-HC-265 80-H-324	260-819	Voyager 1 116,000,000 miles from Saturn
80-HC-538 80-H-688		Voyager 1 76,000,000 miles from Saturn
80-HC-545		Voyager 1 70,000,000 miles from Saturn
80-HC-546 80-H-694	P-22830	Voyager 1 66,000,000 miles from Saturn
80-HC-576 80-H-753	P-22877	Voyager 1 50,000,000 miles from Saturn
80-H-800	P-22974 S-1-6	Saturn's Rings
80-H-801	P-22973 S-1-5	Four photos of Saturn's Rings
80-HC-580	P-22877	Voyager 1 46,000,000 miles from Saturn
80-HC-613 80-H-802	P-22990 S-1-7	Rings of Saturn
80-HC-614		Voyager 1 32,000,000 miles from Saturn
80-HC-615 80-H-804	P-22993 S-1-8	Voyager 1 21,000,000 miles from Saturn
80-HC-616 80-H-805	P-22994 S-1-9	Color enhanced picture of Saturn
80-HC-626 80-H-816	P-23057 S-1-12	Saturn's Moon Titan
80-HC-627 80-H-817	P-23058 S-1-13	Saturn's Moon Tethys
80-H-818	260-906 S-1-10	Saturn's Rings and two small satellites discovered




# Table of Contents (continued)

HQ #	JPL#	Description
80-H-819	P-23053 S-1-11	Spokelike features in Saturn's Rings
80-H-820	P-23059 S-1-14	Saturn's Rings
80-H-823	P-23064 S-1-17	Saturn's Rings
80-H-824	P-23068 S-1-18	Saturn's Rings
80-H-825	P-23069 S-1-19	Saturn's Moon Iapetus
80-H-826	P-23070 S-1-21	Saturn's 15th Moon
80-H-827	P-23075 S-1-24	Cassini Division
80-HC-630 80-H-828	P-23062 S-1-15	Enhanced color of Saturn's Northern Hemisphere
80-HC-631 80-H-829	P-23063 S-1-16	Enhanced color of Saturn's Southern Hemisphere
80-HC-632 80-H-830	P-23067 S-1-20	Enhanced color of Saturn's Southern Hemisphere
80-HC-633 80-H-831	P-23073 S-1-22	Surface of Saturn
80-HC-634 80-H-832	P-23074 S-1-23	Saturn's Rings
80-H-833	260-958 S-1-27	Saturn's F-Ring
80-HC-635 80-H-834	P-23077 S-1-26	Saturn and its Moons Tethys, Enceladus, and Mimas
80-HC-636 80-H-835	P-23076 S-1-25	Saturn's Moon Titan



# Table of Contents (continued)

HQ#	JPL#	Description
80-H-836	P-23082 S-1-29	Saturn's Moon Rhea
80-H-837	P-23083 S-1-30	Saturn's C-Ring
80-H-838	P-23084 S-1-31	Saturn's Moon Tethys
80-H-839	P-23085 S-1-32	Saturn's Moon Rhea
80-H-840	P-23078 S-1-28	Saturn's Moon Dione
80-H-841	P-23099 S-1-39	Saturn's F-Ring
80-H-842	P-23098 S-1-38	Saturn's Moon Mimas
80-H-843	P-23096 S-1-36	Saturn's C-Ring
80-H-844	P-23094 S-1-34	Saturn's Moon Dione
80-H-845	P-23095 S-1-35	Saturn's Moon Tethys



HQ # 80-HC-90 80-H-117

This photograph of Saturn was taken by NASA's Voyager 1 on March 20 at a distance of 312,270,000 kilometers (194,400,000 miles). The picture was computer-enhanced to bring out the rings and other features. One of Saturn's moons, Rhea, can be seen at lower left. Voyager 1 will make its closest approach to Saturn on November 12, 1980; Voyager 2, a sister craft, will fly past in August 1981. The project is directed by the Jet Propulsion Laboratory, Pasadena, California, a government-owned facility managed for NASA by the California Institute of Technology.






FOR RELEASE: PHOTO NO. 3-24-80 PH 80-HC-90 80-H-117

This photograph is a government publication—not subject to copyright.


It may not be used to state or imply the endorsement by NASA or by any EASA employee of a commercial product, process or service, or used in any other manner that might mislead. Accordingly, it is requested that if this photograph is used in advertising, and other commercial fromotion, layout and copy be submitted to NASA prior to release.


This photograph of Saturn was taken by NASA's Voyager 1 on Harch 20 at a distance of 312,270.000 kilometers (194,400,000 miles). The picture was computer-enhanced to bring out the rings and other features. One of Saturn's moon, Rhea, can be seen at lower left. Voyager 1 will make closed approach to Saturn on November 12, 1980; Voyager 2, a sister craft, will fly pas: in August 1981. The project is direct by the Jet craft, will fly pas: in August 1981. The project is direct by the Jet propulsion Laboratory, Pasadena, Calif., a government-owned facility managed for NASA by the California Institute of Technology.



HQ # 80-HC-265 80-H-324 JPL # 260-819


This picture of Saturn taken by NASA's Voyager 1 on June 24, 1980, slightly exceeds the resolution of Earth-based photography of the planet. The picture was taken from a distance of 187,000,000 kilometers (116,000,000 miles) 20 weeks before its November 12 closest encounter. The photograph resolves features 3,500 kilometers (2,175 miles) in diameter, compared to some of the better photographs of Saturn taken from Earth that resolved features with a diameter of 5,000 kilometers (3,100 miles). The Sun crossed Saturn's ring plane last March 20, and has been moving steadily northward since then, providing a steady increase in ring brightness. When this picture was taken, however, the Sun was only 1.9 degrees above the ring plane, and the rings were still unusually dim compared with their usual appearance from Earth. In this photograph, parts of the rings have been digitally brightened, causing the discontinuity in apparent brightness where they crossed Saturn's disk. Tethys, one of Saturn's 11 moons, can be seen below and to the right of the disk of Saturn. The oval just above the rings is caused by removal of a calibration mark on the camera lens. Voyager 1, which flew past Jupiter in March 1979, will have traveled more than two billion kilometers (1.3 billion miles) since its launch from Cape Canaveral, Florida, on September 1, 1977. A sister craft, Voyager 2, is scheduled to encounter the ringed planet on August 27, 1981. The Voyager project is managed for NASA's Office of Space Science by the Jet Propulsion Laboratory, Pasadena, California.







HQ # 80-HC-538 80-H-688

This color photograph of Saturn was produced from black and white images shuttered by NASA Voyager 1 through several different filters. The images were taken August 11, 1980 from a range of 123 million kilometers (76 million miles). The resolution of this photograph far exceeds the best possible Earth-based capability. Belts and zones in the southern hemisphere, though not apparent in this color photo, are readily discernible in the higher contrast black and white images. This is the first color composite in which Encke's Division in Ring A can be seen.







FOR RELEASE:

3-22-33 36- C-538 80--538


The photogoigh is a polenament publication --- not subject to Copy out of .


to may not be used to such or imply the endorsement by 1925th or ky the 1985K employer of a commercial product, process or service or used in any other minor that might midead. Accordingly, it is to quested that it is photograph is used in a feeting, and other continuous promocion, layout and copy by such a section to MASA and to use se.

#### VOY-CER ! PHOTOGRAPHS SATORN

This color photograph of ratum was produced from black and white imager shuftered by NASA Voyager I through several different filters. The image were taken August I. 1980 from a range of 123 million kilometer: (76 million miles). The resolution of this protograph for exceed, the best possible Earth based republify. Belts and zones in the southern hemisphere, though not apparent in this color photo, are readily essertials in the higher contrast black and white images. This is the first color composite in which lackers bivestor in Ring 3 can be seen.

<sup>2-</sup>CID CARDITONICSA of Net 6-11 Amoustaics and Space Administration






FOR RELEASE: 8-28-80 PHOTO NO. 80-HC-545

This photograph is a government publication—not subject to copy right.

It may not be used to state or imply the endocement by NASA or by any NASA employee of a commercial product, process or service, or used in any other manner that might raislead. Accordingly, it is requested that if this photograph is used in advertising, and other commercial promotion, layout and copy be submitted to NASA prior to release.

This color composite is composed of three narrow-angle camera images taken by Voyager I on August 18, 1980 (PDT). At the time these photographs were taken, the spacecraft was 115 million kilometers from Saturn; thus, the smallest resolved features were about 2100 kilometers across. In this composite, the two largest gaps in the rings, Cassini's and Encke's, are seen (Encke's is the easiest to see just on the outermost tips of the rings). The dark cloud belt running across the northern hemisphere is the North Equatorial Belt (NEB). Several frames taken through an ultraviolet filter near this time shown a bright atmospheric feature just north of the NEB, though the feature is not visible in this composite.

These frames represent some of the last taken by Voyager 1 prior to regular, daily Saturn imagining which begins on August 23, 1980 and centinues through November, 1980.



HQ # 80-HC-546 80-H-694 JPL # P-22830

Voyager 1's encounter with the planet Saturn has begun with a series of photos, among them this one of Saturn and three of its satellites. The picture was taken August 24, 1980, 80 days before closest approach, when Voyager 1 was 106,250,000 kilometers (66 million miles) away. A series of dark and light cloud bands appears through high-altitude atmospheric haze in the northern hemisphere. Considerable structure can be seen in the rings: Cassini's Division, between the Considerable structure can be seen in the rings: Cassini's Division, between the A-ring and B-ring, is readily visible. The shadow of the rings on the planet's disk can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The three satellites visible are (left to right) Enceladus (off left can also be seen. The northern hemisphere. The northern hemisphe



## 05907 2 70 CORENGE H

108 RELEASEMIQUSE 26, 1980 10010 MOBJ-HC-546 10010 MOBJ-HC-546




the property is a government publication not subject to copy

¦<sub>e</sub>ųŠ J. C

CHARLE SECURIORS AROUS SECURITY BUT COOK OF MORE SECURITY OF DESCRIPTION OF SECURITY OF DESCRIPTION OF SECURITY OF

· Kungkuoder satsyriding by the grap was the money statistic heavis. MIT 1880 to suitable at most a malane capability and an equality of nach bas mudas happeoins a serio reported that hope of the attents augh medite istes and on at little aspayer, annough theoret tooks "C961 TOT TANK TESS" ATA STO STORE THE FOR PROPER SOLVEN SEEL OF \$125 TO STORE THE THERE THE THREE TERMINER THE CENTER TO DECIDE TO THE TERMINER STATE OF THE TRUMBERS. "Buy uniaq lagunorus upraes , costavas capata, co aba sultipo te sistem pire (Tanalq and mulau Jave encid August To aute and the substance of the substance (วูบัธิเม วัว รูเก.) จิ.ส ส์เกาการ รลัว กูโลเซร สลัมกา หมูก กษตร ลำ รัก ล้ายสา asib a sensig ses that whe to work and to work and the planet a team sk dine gaith-s, and deshadar cartetall eclatered, been himse at east ha เขา อันทรากมาร์ อุเกียนอกรรยดา กามลากิจรู้เอ็ต นะอะท่านดา หนุก การ พิทศ กรุ่นอยประชาก A tews (seling notation as energy being and the ment assets ment typecadde asas ser earled the te to the same researe appeals. and testillates see he sends post moused to any sind with groups rectoring uo salias a nate engan are वसायात व राज्यात काल सामार प्रकार करानावास है। याम्बर्धान




HQ # 80-HC-576 80-H-753 JPL # P-22877

Voyager 1 took this picture of Saturn on September 12, 1980, about nine weeks before its closest approach to Saturn. At that time, Voyager 1 was 81 million kilometers (50 million miles) from the ringed planet and approaching at a velocity of about 1,330,000 kilometers (824,600 miles) per day.

In early March 1980, Saturn was at the point in its orbit where the Sun was in the plane of Saturn's rings; and since then the Sun has been slowly "rising" on the northern side of the rings. At the current illumination angle of 30°, the rings appear much darker than the planet itself, quite unlike most Earth-based photographs.

Very obvious in this image are numerous bands in Saturn's atmosphere, the Cassini and Encke Divisions in the rings, and the ring's shadow on the planet. The Cassini Division is the more prominent gap in the ring's brightness, and the Encke is the fainter gap near the ring edge.



Resolution, The Control

tional sand the sand

FOR RELEASE:

Sept. 17, 1980

MOTO NO.

80-H-753 80-HC-576

This photograph is a government publication-not subject to copy

right.


It may not be used to state or imply the endorsement by NASA or by any NASA employers of a commercial product, process or service or used in any other manner that might misted, Accordingly, it is requested that if this photograph is used in advertising, and other commercial promotion, layout and popy be submitted to NASA prior to release.

## View from Voyager

Voyager 1 took this picture of Saturn on September 12, 1980, about nine weeks before its closest approach to Saturn. At that time, Voyager was 81 million kilometers (50 million miles) from the ringed planet and approaching at a velocity of about 1,330,000 kilometers (824,600 miles) per day.

In early March 1980, Saturn was at the point in its orbit where the Sur was in the plane of Saturn's rings; and since then the sun has been slowly "rising" on the northern side of the rings. At the current illumination angle of 30, the rings appear much darker than the planet itself, quite writke most Earth-based photographs.

Very obvious in this image are numerous bands in Saturn's atmosphere, the Cassini and Encke Divisions in the rings, and the ring's shadow on the planet. The Cassini Division is the more prominent gap in the ring's brightness, and the Encke is the fainter gap near the ring edge.



HQ # 80-H-800 JPL # P-22974 S-1-6

New features in the rings of Saturn that have never been seen before appear in this photo taken by Voyager 1 on October 5, from a distance of 51 million kilometers (32 million miles). The photo has been computer-enhanced to bring out faint details in Saturn's rings. This and similar Voyager photos are the first pictures to show irregular patterns in the rings. Visible in this photo of the rings is a pattern of a dark, fingerlike area that rotates around the planet like a spoke in a wheel. Studies of this and similar photos reveal many similar objects, some that retain their identities for several hours, despite the fact that at the inner edge of the new features, ring objects orbit Saturn once in 9 1/2 hours, while particles at the outer edge take more than an hour longer. Therefore, spokelike features like this should be erased as the inner objects race ahead of the outer ones. But some have been observed that last three or more hours. Voyager imaging-team scientists have not yet solved the question of how the spokes develop or why they remain for hours. It is unlikely, according to the scientists, that the new features are composed of dark objects. Rather, they are more likely to be regions where there are fewer objects, reflecting less light, than other parts of the rings. (Small, square, smudged areas are reseau marks engraved on the camera, and not features of Saturn or its rings.)

неаdquarters Washington, D.C.

#COTO NO. VOYAGET 5-2-6 P-22974 B/V PHOTO NO. VOYAGET 5-2-6 AND PALLY FORK SET SOFTE DEN

rebuta aus ±0


where of their in the authority is a distributed by the contraction of the contraction of

2.2

13 A2AH vot memerodore sett vicerne as meter of base and son year of a sevelous to memero in the sevelous should produce a to sevelous A2AH vos vot vos of the vicerne memoral indicates the memero when it besu to the vicerne and the produce of the


where there are fewer objects, reflecting less light, then other parts are composed of dark objects. Rather, they are more likely to be regions hours, it is unlikely, according to the scientistic, that the new features solved the question of how the spokes develop or why they remem for shead of the outer ones. Voyager imaging-team serentists have not yet spokelike features like this should be erased as the inner objects race while particles at the outer edge take more than an hour longer. Incretors. edge of the new features, ring objects orbit Saturm once in 9 1/2 hours, TOWN SALT to test the tast end endered agreed thrower not sentimed stand mister tedt enoz , achotde taliniz ynam faeven actump talimiz bna pidt seres that rotates eround the planet like a spoke in a wheel Studies of rings. Visible in this photo of the rings is a pattern of a dark. Fingerilke Moyager photos are any first produced of perulant gard and eas solved rapsyov elaffmis bas aid again a countain at stistab datas and paind of becasains of 51 million kilometers (32 million miles). The photo has been computerappear in this photo taken by MASA's Voyager I on Oct. S. from a distance erored need you on the distribution of sent and meren become sent and sentimes well

(Small, square smudged areas are reseau marks engraved on the camera.



HQ # 80-H-801 JPL # P-22973 S-1-5


New features in the rings of Saturn that have never been seen before appear in these four photos taken by Voyager 1 on October 4 and 5, from a distance of 51 million kilometers (32 million miles). The photos have been computer-enhanced to bring out faint details in Saturn's rings. These are the first photos to show irregular patterns in the rings. Visible in these photos of the rings are patterns of dark, fingerlike areas that rotate around the planet like spokes in a wheel. Studies of the photos reveal that some retain their identities for several hours, despite the fact that at the inner edge of the new features, ring objects orbit Saturn once in 9 1/2 hours, while particles at the outer edge take more than an hour longer. Therefore, spokelike features like these should be erased as the inner objects race ahead of the outer ones. But some have been observed that last three or more hours. Voyager imaging-team scientists have not yet solved the question of how these spokes develop or why they remain for hours. It is unlikely, according to the scientists, that the new features are composed of dark objects. Rather, they are more likely to be regions where there are fewer objects, reflecting less light, than other parts of the rings. (Small, square, smudged areas are reseau marks engraved on the camera, and not features of Saturn or its rings.)




NEB, although the feature is not visible in this composite.

These frames represent some of the last taken by Voyager 1 prior to regular, daily Saturn imaging which begins on August 23, 1980 and continues through November 1980.


hemisphere is the North Equatorial Belt (NEB). Several frames taken through an ultraviolet filter near this time show a bright atmospheric feature just north of the





HQ # 80-HC-580 IPL # P-22877

NASA's Voyager 1 took this picture on September 17, 1980, when the spacecraft was still 76 million kilometers (47 million miles) and eight weeks away from its November 12 encounter with the giant ringed planet, capturing five of the planet's moons in the frame. Saturn's largest moon, Titan (considerably larger than Earth's moon), is clearly seen in the upper right corner. The smaller satellites, Mimas and Enceladus, appear to the lower right of the planet, Mimas being the closer one to Saturn. The bright object to the left of the rings is not a moon but an artifact of processing. Voyager 1 will fly past Saturn at a distance of 124,200 kilometers (77,176 miles) on November 12, 1980. This photo is one of a series that began August 24. More than 17,000 photos will be taken of Saturn and its rings and its satellites by Voyager 1. The Voyager project is managed for NASA by the Jet Propulsion Laboratory.




-4 Land double because and land a contract of the contract of the

FOR RELEASE: Sapr. 20, 19-0 PHOTO NO. \$4-4-257 E0-90-580

This photogerph is a queensient publication includeet to dony


It may not be used to state or largery the empoymentally ideas on by my NASA emolorer at a commonwist product, process or service, or used to any other majors that might maken. Accordingly to a consisted that of this proposition used as advertising, and other commercial promotion, by out and copy by administ to MASA prior to extense.

Affairs Regager 1 took this picture on Sapremoer 17, 1401, when the spicement was still to million kilometers (d) collian miles, and dight needs away from its down 12 encounter with the giant chaged planet, appuring the of the planet's moons is the frame. Saturd's large toom, Time toomsiderably larger than baron's coom) is clearly seen in the upper sight corner. The smaller sate intes, there and Tethys, are shown in the upper left corner (apper and lover, respectively). Two of the planet, these being the closer one to sature. The bright object to the fall of the rings is not a moon but an artifact of precassing, as you get I will fly past Sature at a distance of 124, but at one were larger may at the fall than than II, will protoc us one of a series that begin may at the sacrification than II, will protoc will be taken of Sature its comparison taken as sacrification. The Valuager project is managed the MASA to det ampulgion capacities.



HQ # 80-HC-613 80-H-802 JPL # P-22990 S-1-7

This computer composite of four Voyager 1 photos contains a wealth of new information on the rings of Saturn, including subtle color variations and new structural features. Voyager 1 took the pictures on October 13, 1980, while 40 million kilometers (25 million miles) from the planet. They were combined and enhanced in the Image Processing Lab at Jet Propulsion Laboratory to make this false-color picture. Although colors are false (a product of the computerenhancement process), color variations are real. Satellite Mimas is at bottom edge of photo; bright spot at right edge is an artifact of processing. Portions of four of Saturn's rings can be seen: small arc-shaped feature at lower right is part of the F-ring, discovered by Pioneer 11 during its Saturn encounter in 1979. Next ring inward is the A-ring, split by the dark Encke's Division. Dark region between A-ring and B-ring is Cassini's Division, filled with material discovered by Voyager 1. Considerable variations in distribution and brightness of material can be seen in the B-ring. Innermost ring visible here is the C-ring, which also shows variations in distribution and brightness of material. Color variations can be seen in the planet itself.



-


ale of the least o

108 18 E-Skickuper rd, 1880 14070 NBU-BC-563 80-H-803 Yoyager S-1-7 P-229900


The photograph is a government publication inno subject to coperate

It may not be used to state or analy the endorsement by NASA or by any MASA employed of a commercial product, process or service, or used in any other manner that might mushed. Accordingly, it is requested that if this photograph is used in acvertising, and other commercial promotion, Leyous and copy be submitted to NASA arior to release

This computer composite of four Voyager 1 photos contains a wealth of men information of the rings of Saturn, including subtle color variations and now structural features. Voyager 1 took the pictures on October 13, 1980, while 40 million kilometers (25 million miles) from the planet. They were combined and enhanced in the Image Processing Lab at Jet Propulsion Laboratory to make this false-color picture. Although colors are false (a product of the computer-enhancement process), color variations are rail. Sazellite Mimas is at houton edge of photo; bright spot at right adge is an artifect of processing. Fortions of four of Saturn's rings can be seen. Small erc-shaped feature at lower right is part of the Foring, discovered by Ftoneer Li during its Saturn encounter in 1979. Hext ring inwird is the A-ring, split by the dark Enke Division. Dark region between A-ring and B-ring is Cassini Division, filled with material discovered of Voyager 1. Considerable variations in distribution and brightness of material can be seen in the 8-ring. Innermost ring wisible here is the Caring, which also shows variations in distribution and beightness of material Color variations can be seen in the planet itseld.



The Voyager 1 spacecraft acquired this photograph of Saturn on October 5, 1980, from a distance of 51 million kilometers (32 million miles). At this time Voyager 1 was over 1.5 billion kilometers (940 million miles) from Earth and less than 6 weeks away from its November 12, 1980 closest approach to Saturn. The several spots appearing on the planet's disk are artifacts of this unenhanced photo.







PHOTO: NO. 301 (C.4)

This protograph is a government aublication on 10,3/20110 2007

t may not be used to these or implies to endorsome by MASA or by any NASA employee of a commerce of product, promision which or used to end other manner that height this lead. Accordingly, it is requested that to this photograph is used in advertising, sext other commercial promotion, lawout and copy be submitted to his Se, print to release.

## VINAGER

The Voyager I spacecraft acquired this protograph of Smurn on actiber 5, 1980, from a distance of 53 million allowed to 12 million miles). It this time Voyager I was over 1.5 billion kilometers .940 million this time Voyager I was over 1.5 billion kilometers .940 million in 1981 miles) from Earth and Twee that I was away from this is venue: 2, 1981 miles) from Earth and Twee that I spots topograph on the planet 3 million are arrificals of this mission who photo.



HQ # 80-HC-615 80-H-804 JPL # P-22993 S-1-8

Voyager 1 took this photograph of Saturn on October 18, 1980, when the spacecraft was 34 million kilometers (21.1 million miles) from the planet. The photograph was taken on the last day Saturn and its rings could be captured within a single narrow-angle camera frame as the spacecraft closed in on the planet for its nearest approach on November 12, 1980. Dione, one of Saturn's inner satellites, appears as three color spots just below the planet's south pole. An abundance of previously unseen detail is apparent in the rings. For example: a gap in the dark, innermost ring, called the C-ring, or Crepe ring, is clearly shown; material is seen within the relatively wide Cassini Division, separating the middle, B-ring from the outermost ring, the A-ring. Encke's Division is shown near the outer edge of the A-ring. The detail in the rings' shadows cast on the planet is of particular interest: the broad, dark band near the equator is the shadow of the B-ring; the thinner, brighter line just to the south is the shadow of the less dense A-ring. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.



Special Administration


FIORD NO.

80-40-635 80-4-804 Voyager S-1-8

This photograph is a government publication-not subject to copy right.


It may not be used to state or imply the endorsement by NASA or by any NASA employee of a commercial product, process or service, or used in any other manner that might mislead. Accordingly, it is requested that if this photograph is used in advertising, and other commercial promotion, layout and copy be submitted to NASA prior to release

Voyager 1 took this photograph of Saturn on October 18, 1980, when the spacecraft was 34 million kilometers (21.1 million miles) from the planet. The photograph was taken on the last day Saturn and its rings could be captured within a single narrow-angle camera frame as the spacecraft blosed in on the planet for its nearest approach on November 12, 1980. Dione, one of Saturn's inner satellites, appears as three color spcts just below the planet's south pole. An abundance of previous y unseen detail is apparent in the rings. For example: a gap in the dark, innermost ring, called the C-ring, or Crepe ring, is clearly shown; material is seen within the relatively wide Cassini Division, separating the middle, B-ring from the outermost ring, the A-ing. The Encke Division is shown near the outer edge of the A-ring. The detail in the rings' shadows cast on the planet is of particular interest: the broad. dark band near the equator is the shadow of the B-ring; the thinner, brighter line just to the south is the shadow of the less dense A-ring. The Voyager Project is managed for NASA by the Jet Propulsion Laboratory. Pasadena. California.



HQ # 80-HC-616 80-H-805 JPL # P-22994 S-1-9


This image of Saturn, taken by the Voyager 1 spacecraft on October 18, 1980, was color-enhanced to increase the visibility of large, bright features in Saturn's North Temperate Belt. It is believed that these spots might closely resemble gigantic convective storms (similar to, but much larger than, thunderstorms in Earth's atmosphere) with upwelling from deep within Saturn's atmosphere. The nature of the dark spots like the one visible on the northern edge of the belt is not yet clearly understood, though they seem to resemble equally mysterious features seen in Jupiter. The largest violet-colored cloud belt (its true color is brownish) is Saturn's North Equatorial Belt. The distinct color difference between this and other belts and zones may be due to a thicker haze layer covering the northern portion of the belt. The southern hemisphere of the planet (below the rings) appears bluer than the northern hemisphere because of the increased scattering of sunlight upon that area due to the spacecraft's point of view. Numerous gaps and divisions in the rings are becoming more apparent as Voyager 1 approaches Saturn. The newly discovered dark radial features in the rings are faintly shown on the lower right portion of the rings. Three separate Voyager 1 images taken through ultraviolet, green, and violet filters were used to construct this blue, green, and red color composite of Saturn. The lower edge of the rings was "clipped" due to a slight drift of the spacecraft. Color spots in the rings are an artifact of image processing. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.



on the state of th


The protograph is a second without the property of contributions and property of the protograph of the property of the protograph of the p

o frequency galaxies, had not by the contragers of the contract on their I on a minum model of minusely one of the confidence while w. in litury a placed force error surper of the cultivers hat the seor soft tips o research country and tribe age. I will at the to and a more chain and experience compared and another comelling company of the document and construction and company the documents of the se se visi le a line softhere each to be 1835 a per let liest. a governor, nouge they ago to relace a color onlyster the leatures en process par l'impliment de la contra del contra de la contra del many common quarter of the assimultaness. The state of the s This wild thought the control of the large of the late a three t fi / co nower with an error view of the contracts postable of the partie. The cathers was notes to the reason (be on the rings of size or bluer to be doubtern are compared to the country of the approved to the approved to the contact in the compression of the property of the contract of the cont ក្រុម ខេត្ត ខេត្តបន្ទាប់ ការី ឆ្នាំ ១០១៩ ខែ១០១៩ ខេត្ត ១០១៩ ខេត្ត ការប្រជាធិប្បធានការប្រជាធិប្បធានការប្រជាធិប្ប ខេត្តបន្ត ខេត្តបន្ទាប់ ការី ខេត្តខេត្តបន្ទាប់ ខេត្ត ខេត្តបន្ទាប់ ខេត្តបន្ទាប់ បានប្រជាធិបត្តបានការប្រជាធិបត្តប ins first to a bise, group and sealed to conscious of lacable. The lowe operations sings where slipping due to a single in the or his spacedra? In the page of the single spacedra? I park the product for managed for wells of the method as the car tended materials C3 , 15 ₹5™ , 1



HQ # 80-HC-626 80-H-816 JPL # P-23057 S-1-12


Titan, the largest of Saturn's 14 known satellites, shows little more than the upper layer of clouds covering the moon in this Voyager 1 picture, taken on November 4, 1980 at a range of 12 million kilometers (7,560,000 miles). The orange-colored haze, believed to be composed of photochemically produced hydrocarbons, hides Titan's solid surface from the Voyager cameras. Some weak shadings in the clouds are becoming visible. However, note that the satellite's southern (lower) hemisphere is brighter than the northern. It is not known whether these subtle shadings are on the surface or are due to clouds below a high haze layer. The Voyager project is managed by JPL for NASA.





HQ # 80-HC-627 80-H-817 JPL # P-23058 S-1-13


Saturn and two of its moons, Tethys (above) and Dione, were photographed by Voyager 1 on November 3, 1980, from 13 million kilometers (8 million miles). The shadows of Saturn's three bright rings and Tethys are cast onto the cloud tops. The limb of the planet can be seen easily through the 3,500-kilometer wide (2,170 miles) Cassini Division, which separates ring A from ring B. The view through the much narrower Encke Division, near the outer edge of ring A, is less clear. Beyond the Encke Division (at left) is the faintest of Saturn's three bright rings, the C-ring or Crepe ring, barely visible against the planet. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.



This photograph is a present published memor midject to copy


It may not be used to lesse or leaply the conformation by NASA or by any MASA employs. In estimateful are det, process or servins, or used in any other man or that might misk id. Appointingly, in is equested that if this photo pept is used in adicatising, and other occamercial promotion, layer and copy be soon at ad to NASIA prior to roleasa.


Satern and havef the moons. Tay ye (above) a 1 Dione, were shotographed by Voyager . In November 3, 1930 from 13 million k limeters (8 million males). The stadows of Saturn's three bright lags and Teth, stare east once the near tops. The limb the planet in be seen easily through the S50 will be started (2170 moles) Cassini lyision, which reparates ring A from ing b. The view the aich the much tarrower Encks Division, near the publication of ring A to less clear. Beyond the Encke Distator (at left) is he deintest of Sa mais three beight rings, the Coring or creps roug, bely visible againt the planet. The Joyager Project is straiged by the Car Propulation to protony for AsA.



HQ # 80-H-818 JPL # 260-906 S-1-10


Two small satellites of Saturn recently discovered in images taken by Voyager 1 are seen in this photograph taken by the spacecraft on October 25, 1980, at a distance of 25 million kilometers (15.5 million miles) from Saturn. The smaller, innermost of the two has a radius of about 250 kilometers (155 miles) and is seen near the bottom edge of the A-ring (left). It travels an orbit between the A-ring and the F-ring. (The F-ring is barely visible in the top left corner above the A-ring in this enhanced image.) The second satellite, seen just outside of the F-ring (top), is about 300 kilometers (186 miles) in radius. Scientists believe the dimensions of the F-ring may be determined by these two satellites, which orbit on either side of the ring. Recently discovered concentric features in the rings and material within Cassini's Division are easily visible. Circular black spots in the rings are reseau marks engraved on the camera and not features in the rings. Voyager 1 makes its closest approach to Saturn on November 12, 1980. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-819 JPL # P-23053 S-1-11


Dark spokelike features in Saturn's rings are seen revolving around the planet with the rings' orbital motion in these six photographs taken by NASA's Voyager 1 spacecraft on October 25, 1980. The images were taken in sequence (from upper left to lower right) approximately every 15 minutes at a distance of about 24 million kilometers (14.9 million miles) from the planet. The rotation of the spokelike features, visible in the brightest part of the rings, is recorded in each frame. Because the outer part of the rings revolves more slowly than the inner rings, the differential motion is thought to cause the features to dissipate. However, the radial features are apparently ubiquitous and are regenerated by some unexplained mechanism. Dark round spots on the rings and planet are reseau marks engraved on the camera and are not features of Saturn. Voyager 1 makes its closest approach on November 12, 1980. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.



HQ # 80-H-820 JPL # P-23059 S-1-14

This photograph of Saturn's rings taken by NASA's Voyager 1 on November 3, 1980 at a range of 13 million kilometers (8.19 million miles) shows complex structure within the Cassini Division (separating the A- and B-rings) never seen before. Seen within the division are four bright ringlets separated by darker components and an outer ringlet of medium brightness. All of these internal features are bounded by an inner and outer dark division. Such structure within the Cassini Division suggests that some mechanism other than the interaction between Saturn's satellites and the orbit of the ring particles must be responsible for the detailed structure of Saturn's rings. The Cassini Division, the widest in Saturn's rings, was first seen by Jean Dominique Cassini in 1675. The Encke Division was first seen by Johann Franz Encke in 1837, and appears near the outer edge of the A-ring (the outermost ring seen here). The Voyager project is managed for NASA by the Jet Propulsion Laboratory.






HQ # 80-H-823 JPL # P-23064 S-1-17

This two-image mosaic of Saturn's rings, taken by NASA's Voyager 1 on November 6, 1980 at a range of 8 million kilometers (4.9 million miles), shows approximately 95 individual concentric features in the rings. The extraordinarily complex structure of the rings is easily seen across the entire span of the ring system. The ring structure, once thought to be produced by the gravitational interaction between Saturn's satellites and the orbit of ring particles, has now been found to be too complex for this explanation alone. The 14th satellite of Saturn, discovered by Voyager, is seen (upper left) just outside the narrow F-ring, which is less than 150 kilometers (93.2 miles) wide. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.

Headquarters Washington, D.C.




w ≱ger of the control of the contro


以上の1975年 1997年 - 19

> ner Name – Name – Name

. . . . . . . . . .

				i
				i
			•	
· :				
-				
	•			





Washington, D. L. Co.




FOR RELEASE: Nov. 7, 1980 PHOTO NO. 80-H-823

This obstograph is a government publication—not subject to copy


To may not be used to state or imply the andorsoment by NASA or by any NASA employee of a commercial product, process or service, at use a many other manner that might related. Accordingly, it is remeasure that if this photograph is used in advertising, and other commercial promotion, layout and copy be submitted to NASA prior to release.


Inis two-image massic of Saturn's rings, taken by NASA's Voyager 1 on Nov. 6. 1980 at a range of 8 million kilometers (4.9 million miles). Thoms approximately 95 individual concentric features in the rings. The extraordinarily complex structure of the rings is easily seen across the entire span of the ring system. The ring structure, once thought to be produced by the gravitational interaction between Saturn's satellites and the orbit of ring particles, has now been found to be too complex for this explanation alone. The 14th satellite of Saturn, discovered by Noyager is seen (upper left) just outside the narrow F-ring, which is less than 150 kilometers (93.2 miles) wide. The Voyager Project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, Calif.



HQ # 80-H-826 JPL # P-23070 S-1-21


Voyager 1 has found a 15th moon orbiting Saturn, visible near the bottom of this picture taken on November 6, 1980, when the spacecraft was still 8 million kilometers (5 million miles) from Saturn. Voyager imaging team scientists discovered the moon November 7, 1980 in the first of several programmed searches for new satellites of Saturn. The unique location of the 15th satellite, just 800 kilometers (500 miles) outside the outer edge of the A-ring, is especially significant in that this small body, approximately 100 kilometers (50 miles) in diameter, may be responsible for defining the outer edge of Saturn's bright ring system. The orbital period of the new satellite is approximately 14 hours, 20 minutes, the shortest orbit of any of Saturn's known satellites. The very narrow F-ring, approximately 4,000 kilometers (2,500 miles) outside the outer edge of the A-ring, is seen prominently in this picture. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-827 JPL # P-23075 S-1-24

The Cassini Division in Saturn's rings was photographed by Voyager 1 on November 8, 1980, at a distance of 6 million kilometers (4 million miles). This prominent feature in the ring system, discovered by Cassini in 1675, is defined as the region between the two dark ringlets (1 1/2 inches apart on this print). Within the Cassini Division can now be seen a number of individual features (from its outer boundary to the inner boundary): a medium dark ringlet, 800 kilometers (500 miles) wide; four brighter ringlets, approximately 500 kilometers (300 miles) wide and separated by dark divisions; and a new, barely visible, narrow, bright ringlet at the inner boundary. The width of this newest Cassini ringlet appears to be not more than 100 kilometers (50 miles). The Voyager project is managed by the Jet Propulsion Laboratory for NASA.






HQ # 80-HC-630 80-H-828 JPL # P-23062 S-1-15


This enhanced color image of the northern hemisphere of Saturn taken by NASA's Voyager I on November 5, 1980 at a range of 9 million kilometers (5.5 million miles) shows a variety of features in Saturn's clouds: Small-scale convective cloud features are visible in the brown belt; an isolated convective cloud with a dark ring is seen in the light brown zone; and a longitudinal wave is visible in the light blue region. The smallest features visible in this photograph are 175 kilometers (108.7 miles) across. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-HC-631 80-H-829 JPL # P-23063 S-1-16


This false color image of Saturn's southern hemisphere was taken by NASA's Voyager 1 on November 6, 1980 from a distance of 8 million kilometers (4.9 million miles). The color contrasts are an indication of the divisions between belts and zones in this region. The shadow of the satellite Dione is seen as a dark circle on the face of the planet. A large red cloud feature 12,000 kilometers (7,456 miles) in length is evident in the south polar latitudes (55 degrees south) and has been seen continuously by Voyager since August 1980. The feature is most easily seen in the ultraviolet. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-HC-632 80-H-830 JPL # P-23067 S-1-20

This false color image of Saturn's southern hemisphere taken by NASA's Voyager 1 on November 6, 1980 shows the unique red oval cloud feature located at 55 degrees south latitude. The photograph was taken by the spacecraft at a distance of 8,500,000 kilometers (5,300,000 miles) from Saturn. The difference in color between the red oval and surrounding bluish clouds indicates that material within the oval contains a substance that absorbs more blue and violet light than the bluish clouds. Voyager imaging team scientists first observed the oval in August 1980, and the feature has seemed to retain its appearance since its discovery. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-HC-633 80-H-831 JPL # P-23073 S-1-22


Two brown ovals, at right, some 10,000 kilometers (6,000 miles) across, were found at approximately 40 degrees and 60 degrees latitude in Saturn's northern hemisphere by Voyager 1. The photo was taken on November 7, 1980, from a range of 7,500,000 kilometers (4,600,000 miles). The polar oval (upper left) has a structure similar to the Saturn red spot located in the southern polar latitudes. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.






HQ # 80-HC-634 80-H-832 JPL # P-23074 S-1-23

This computer composite of photos obtained October 30 illustrates the capability of the Voyager narrow-angle camera to portray scenes not visible to the human eye. Normally, at visible wavelengths, Saturn's rings appear quite uniform in color and show dramatic contrast between the bright A- and B-rings and the much fainter C-ring and Cassini Division. Here, in ultraviolet light, the contrast between bright and dark features is greatly reduced. The blue in the photo is derived from an ultraviolet image. Green and red are produced by blue and green filters in the camera system. The colored "spokes" seen faintly in the B-ring result from the time lapse between camera shuttering of the three images. Because the features have moved between exposures, they erroneously show up as color differences. Range from Voyager 1 to Saturn during this photo sequence was about 18 million kilometers (11 million miles). The Voyager project is managed by the Jet Propulsion Laboratory for NASA.




· ·



HQ # 80-H-833 JPL # 260-958 S-1-27


Variations in the brightness of Saturn's thin F-ring are visible in this photograph taken by NASA's Voyager 1 on November 8, 1980 at a range of 7 million kilometers (4 million miles). The planet's 14th satellite, discovered by Voyager 1, is seen just inside the F-ring (bottom). Variations in the brightness of the F-ring, which is less than 100 kilometers (60 miles) wide, may be the result of clumping in the ring material. The features are seen at the top and again near the left edge of the ring in this image. The "gap" in the ring (left center) is not real but is the location of a reseau mark engraved on the Voyager cameras. These same bright features in the F-ring have been observed for several days and appear to move at the orbital rate of the ring particles. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-HC-635 80-H-834 JPL # P-23077 S-1-26


Saturn and its satellites Tethys (outer left), Enceladus (inner left), and Mimas (right of rings) are seen in this mosaic of images taken by NASA's Voyager 1 on October 30, 1980 from a distance of 18 million kilometers (11 million miles). The soft, velvety appearance of the low-contrast banded structure and increased reflection of blue light near the perimeter of the Saturn disk are due to scattering by a haze layer above the planet's cloud deck. Features larger than 350 kilometers (220 miles) are visible. The projected width of the rings at the center of the disk is 10,000 kilometers (6,000 miles), which provides a scale for estimating feature sizes on the image.






HQ # 80-HC-636 80-H-835 JPL # P-23076 S-1-25

The clouds covering Saturn's satellite Titan are seen in their true colors in this image taken by Voyager 1 on November 9, 1980 at a distance of 4.5 million kilometers (2.8 million miles). Titan's southern hemisphere appears as the lighter region with a well-defined boundary between the clouds around the satellite's equator. The northern polar region appears darker than clouds in the neighboring area. The terminator of Titan is at the right; darkening in this area is consistent with extensive cloud layers covering the satellite. Titan's surface is believed to be shrouded by aerosols rather than the convective clouds found in the atmospheres of Jupiter and Saturn. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-836 JPL # P-23082 S-1-29

Broad areas of light and dark material seen in this enhanced Voyager 1 picture of Saturn's satellite Rhea suggest varying amounts of fresh ice on the moon's surface. A few small bright spots are visible and are similar in appearance to small fresh impact craters seen in Voyager images of the Galilean satellites of Jupiter taken at this resolution. The photograph, taken November 10, 1980 at a range of 2.7 million kilometers (1.6 million miles), shows features about 50 kilometers (30 miles) in scale. Near the terminator (lower right) are several features which indicate topographic relief. The two circular features may be large impact craters perhaps a hundred kilometers (60 miles) across. (The extreme brightness of the southern (lower left) region is probably not real and is a result of image processing.) Rhea is one of the larger of Saturn's inner satellites with a diameter of about 1,500 kilometers (900 miles) and is known to be covered at least partially with water frost. It may be mostly icy in nature. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-837 JPL # P-23083 S-1-30


Both the limb of Saturn and the shadow of its ring system are seen through the transparent C-ring in this striking picture taken by NASA's Voyager 1 on November 9, 1980 at a distance of 4.5 million kilometers (3 million miles). Gaps and regions of high transparency are seen throughout the C-ring, especially in the area closest to the opaque B-ring. Shadows of the A-, B-, and C-rings are clearly visible on the disk of Saturn. The C-ring shadow showing the gaps described above is the uppermost shadow; below this is the very black shadow of the opaque B-ring; then the Cassini Division; and at the bottom, the shadow of the A-ring.






HQ # 80-H-838 JPL # P-23084 S-1-31


A large circular feature, which may represent either an albedo feature or a hill, is seen in this Voyager 1 picture of Saturn's satellite Tethys taken November 11, 1980 at 4:47 a.m. (PST) from a range of approximately 2 million kilometers (1.2 million miles). The feature, visible near the center of the satellite, is about 180 kilometers (100 miles) across. The terminator is seen at the extreme right edge of the satellite, so the dark areas on the left side of Tethys result from darker surface material. The smallest surface detail that can be seen in this image is about 36 kilometers (22 miles) across. Tethys has a diameter of about 1,000 kilometers (600 miles), which is about one-third the size of Earth's Moon. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-839 JPL # P-23085 S-1-32

Bright streaks and blotches are visible against a darker background on the surface of Saturn's satellite Rhea, seen in this Voyager 1 image taken November 11, 1980 from a range of 1,925,000 kilometers (1,196,000 miles). Even the dark areas, thought to be water frost and ice, are fairly bright with about 50 percent reflectance. The bright streaks may be related to impacts by objects that throw out pulverized ice grains from beneath the ice-covered surface. Some of the bright streaks are not straight but have a curved appearance similar to the grooved, icy terrain on Jupiter's satellite Ganymede seen in Voyager photographs taken at this resolution. Scientists do not yet know if a satellite of Rhea's size (approximately 1,500 kilometers or 900 miles in diameter) can have an active thermal history like Ganymede's, but higher resolution photographs taken by Voyager should reveal clues to its history. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.






HQ # 80-H-840 JPL # P-23078 S-1-28


This picture of Saturn's satellite Dione was taken by NASA's Voyager 1 on November 9, 1980 from a distance of about 4.2 million kilometers (2.6 million miles). Light and dark patches are visible on the moon's surface, reminiscent of features seen on Jupiter's satellite Ganymede during Voyager's Jupiter encounter last year. The bright spots may be rays emanating from impact craters on Dione's surface. Dione is about 1,110 kilometers in diameter, about one-third the size of Earth's moon. The smallest detail seen in this image is about 78 kilometers (48 miles) across. The Voyager project is managed for NASA by the Jet Propulsion Laboratory, Pasadena, California.






HQ # 80-H-841 JPL # P-23099 S-1-39


Saturn's F, or outermost, ring was photographed from the unilluminated face of the rings by Voyager 1 at a range of 750,000 kilometers (470,000 miles). Complex structure is evident, with several components seen. Two narrow, braided, bright rings that trace distinct orbits are evident. Visible is a broader, very diffuse component about 35 kilometers (20 miles) in width. Also seen are "knots," which probably are local clumps of ring material, but may be mini-moons. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.






HQ # 80-H-842 JPL # P-23098 S-1-38

This Voyager 1 picture of Mimas shows a large impact structure at 110° W Long., located on that face of the moon which leads Mimas in its orbit. The feature, about 130 kilometers in diameter (80 miles), is more than one-quarter the diameter of the entire moon. This is a particularly interesting feature in view of its large diameter compared with the size of the satellite, and may have the largest crater diameter/satellite diameter ratio in the solar system. The crater has a raised rim and central peak, typical of large impact structures on terrestrial planets. Additional smaller craters, 15-45 kilometers in diameter, can be seen scattered across the surface, particularly along the terminator. This photo was taken on November 12 at 5:05 a.m. PST, from a range of approximately 660,000 kilometers (400,000 miles). Mimas is one of the smaller Saturnian satellites with a low density implying its chief component is ice. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.







HQ # 80-H-843 JPL # P-23096 S-1-36

High resolution detail in Satum's C-ring is illustrated in this composite photo obtained November 10 by Voyager 1 from a range of 3 million kilometers (2 million miles). The horizontal line through the center marks the border between the two photos; at top the trailing ansa of the rings, and bottom the leading ansa. The dark gap in the center of both pictures shows the bright ring—narrowed in the lower picture and slightly broadened and displaced within the gap in the upper picture. This illustrates the eccentricity of the C-ring. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.


National Aeronautics and Space Administration






HQ # 80-H-844 JPL # P-23094 S-1-34

Circular impact craters up to about 100 kilometers (60 miles) in diameter are seen in this view of Saturn's icy moon Dione. The image was taken by Voyager 1 from a range of 790,000 kilometers (500,000 miles) at 2:20 a.m. PST on November 12. Bright, wispy markings form complex arcuate patterns on the surface. These markings are slightly brighter than the brightest features seen by Voyager on Jupiter's moons, suggesting that they are surface frost deposits. The patterns of the bright bands hint at an origin due to internal geologic activity, but the resolution is not yet sufficient to prove or disprove this idea. Dione's diameter is only 1100 kilometers (700 miles), much smaller than any of Jupiter's icy moons. It thus belongs to a class of small, icy objects never observed before the Voyager 1 Saturn encounter. The view here is of the face which trails in orbit. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.





HQ # 80-H-845 JPL # P-23095 S-1-35

The heavily cratered surface of Tethys was photographed at 1:35 a.m. PST on November 12 from a distance of 1.2 million kilometers (750,000 miles) by Voyager 1. This face of Tethys looks toward Saturn and shows a large valley about 750 kilometers long and 60 kilometers wide (500 by 40 miles). The craters are probably the result of impacts and the valley appears to be a large fracture of unknown origin. The diameter of Tethys is about 1000 kilometers (600 miles) or slightly less than 1/3 the size of our Moon. The smallest feature visible on this picture is about 24 kilometers across. The Voyager project is managed by the Jet Propulsion Laboratory for NASA.

